



Dilated Cardiomyopathy Overview

Ray E Hershberger, MD¹ and Elizabeth Jordan, MS, LGC²

Created: July 27, 2007; Revised: April 7, 2022.

Summary

The purpose of this overview is to increase clinician awareness of the genetic basis of dilated cardiomyopathy (DCM) and the benefits of early diagnosis and management to individuals with genetic DCM.

The following are the goals of this overview.

Goal 1

Define DCM.

Goal 2

Identify the categories of DCM.

Goal 3

Provide the evaluation strategy of a proband with nonsyndromic DCM.

Goal 4

Provide a basic view of genetic risk assessment of at-risk asymptomatic relatives of a proband with DCM to inform cardiac surveillance and allow early detection and treatment of DCM to improve long-term outcome.

1. Dilated Cardiomyopathy (DCM): Definition

The diagnosis of DCM is established when both of the following are present:

- **Left ventricular enlargement.** Enlargement is most commonly assessed in adults by either echocardiography or cardiac MRI. Because of rapid growth in children, expert cardiovascular assessment is recommended to assess left ventricular enlargement in the pediatric population.

Author Affiliations: 1 Professor of Medicine, Divisions of Human Genetics and Cardiovascular Medicine, The Ohio State University, Columbus, Ohio; Email: ray.hershberger@osumc.edu. 2 Assistant Professor, Division of Human Genetics, The Ohio State University, Columbus, Ohio; Email: elizabeth.jordan@osumc.edu.

Copyright © 1993-2024, University of Washington, Seattle. GeneReviews is a registered trademark of the University of Washington, Seattle. All rights reserved.

- **Systolic dysfunction, a reduction in the myocardial force of contraction.** An ejection fraction of less than 50% is considered systolic dysfunction. The left ventricular ejection fraction is the most commonly used clinical measure of systolic function, and is usually estimated from a two-dimensional echocardiogram or from cardiac MRI. Another noninvasive approach is a cardiac nuclear study. Ejection fractions can also be estimated from a left ventricular angiogram.

Note: [Arrhythmogenic right ventricular cardiomyopathy \(ARVC\)](#) with predominant left ventricular involvement may present as DCM [Sen-Chowdhry et al 2008].

DCM usually initially manifests in adults in the fourth to sixth decade, although it may present at any age (prenatally; in infancy, early or late childhood, or adolescence; or in the elderly). Extensive additional clinical and genetic information on DCM is available [Burkett & Hershberger 2005, Sivasankaran et al 2005, Judge 2009, Dellefave & McNally 2010, Hershberger et al 2010a, Jordan & Hershberger 2021].

Persons with DCM may be asymptomatic for a number of years. Manifestations usually occur late in the disease course with one or more of the following findings:

- **Heart failure.** Symptoms include those of congestion (edema, orthopnea, paroxysmal nocturnal dyspnea) and/or reduced cardiac output (fatigue, dyspnea on exertion).
- **Arrhythmias and/or conduction system disease.** These commonly accompany advanced cardiomyopathy and heart failure. Some genetic causes (e.g., pathogenic variants in *DES*, *FLNC*, *LMNA*, and *SCN5A*) may involve prominent conduction system disease or arrhythmias out of proportion to the degree of left ventricular dysfunction.
- **Thromboembolic disease.** Stroke or systemic embolus secondary to left ventricular mural thrombus may also occur.
- **Pregnancy.** Peripartum or pregnancy-associated cardiomyopathy (PPCM/PACM) that occurs during or soon after pregnancy was once considered distinct from DCM, but is now recognized as a part of the clinical spectrum of DCM.

2. Dilated Cardiomyopathy (DCM): Categories

DCM can be categorized as acquired, syndromic, or nonsyndromic (Figure 1).

Acquired (Secondary) DCM


The most common cause of acquired DCM is **ischemic injury**, such as that caused by prior myocardial infarction from coronary artery disease.

Other less common causes include valvular and congenital heart disease, toxins (most commonly, anthracyclines or other chemotherapeutic agents; various drugs with idiosyncratic reactions), thyroid disease, inflammatory or infectious conditions, severe long-standing hypertension, and radiation. While emerging evidence suggests that DCM arising after chemotherapy exposure may also have a genetic background, the clinical relevance of this information is currently undefined [Garcia-Pavia et al 2019].

Note: Acquired DCM will not be discussed further in this overview.

Syndromic DCM

In *GeneReviews*, "syndromic" refers to a disorder characterized by a constellation of phenotypic features that either: (1) specifically suggests the diagnosis (which can be confirmed by molecular genetic testing) or (2) allows diagnosis of the disorder in the absence of confirmatory molecular genetic findings. A selected list of syndromic DCM is provided in Table 1 [Hershberger et al 2009, Hershberger et al 2013].



1. ≥2 close relatives with DCM (including proband)
2. Based on American College of Medical Genetics and Genomics (ACMG)/Association for Molecular Pathology (AMP) criteria

Figure 1. Categories of dilated cardiomyopathy

Note: Syndromic DCM will not be discussed further in this overview.

Table 1. Selected List of Syndromic Dilated Cardiomyopathy

Disorder ¹	Gene(s)	MOI	Other Clinical Features	Comments
Barth syndrome	<i>TAFAZZIN (TAZ)</i>	XL	<ul style="list-style-type: none"> Neutropenia Muscle weakness Growth delay Infantile/early-childhood onset 	
Carvajal syndrome (OMIM 605676)	<i>DSP</i>	AR	<ul style="list-style-type: none"> Woolly hair Palmoplantar keratoderma 	
Duchenne & Becker muscular dystrophy	<i>DMD</i>	XL	In males: <ul style="list-style-type: none"> Muscle weakness ↑ serum CK levels Loss of ambulation in childhood or later in life 	Heterozygous females may present w/isolated DCM.
Emery-Dreifuss muscular dystrophy	<i>EMD</i> <i>FHL1</i>	XL	<ul style="list-style-type: none"> Joint contractures ↑ serum creatine kinase (CK) levels Childhood- or adult-onset muscle weakness 	Conduction system disease &/or arrhythmias are common.
	<i>LMNA</i>	AD AR		

Table 1. continued from previous page.

Disorder ¹	Gene(s)	MOI	Other Clinical Features	Comments
HFE hemochromatosis	<i>HFE</i>	AR	<ul style="list-style-type: none"> • Cirrhosis • Diabetes • Hypermelanotic pigmentation • ↑ serum iron & ferritin levels 	Nondilated &/or infiltrative cardiomyopathy is more frequent than DCM.
Laing distal myopathy	<i>MYH7</i>	AD	<ul style="list-style-type: none"> • Facial weakness • Childhood-onset weakness of ankles, great toes, finger extensors, & neck flexors 	
Mitochondrial DCM (See Mitochondrial Disorders Overview .)	mtDNA	Mat	Complex phenotypes incl: <ul style="list-style-type: none"> • Focal segmental glomerulosclerosis • Kearns-Sayre syndrome 	

AD = autosomal dominant; AR = autosomal recessive; Mat = maternal inheritance; MOI = mode of inheritance; XL = X-linked
1. Disorders are in alphabetic order.

Nonsyndromic DCM

Individuals with DCM who do not have acquired (secondary) DCM or syndromic DCM (Table 1) have nonsyndromic dilated cardiomyopathy (defined for this *GeneReview* as DCM with no other systemic involvement). See Table 2 for a current list of known DCM-related genes organized by strength of ClinGen classification and alphabetically [Jordan et al 2021].

The [Clinical Genome Resource \(ClinGen\) DCM Gene Curation Expert Panel](#) has classified DCM-related genes using the ClinGen framework for the strength of their relationship with monogenic, nonsyndromic DCM. A summary of the data curated for each gene can be accessed at [ClinGen Gene Validity Classification](#).

Note: Left ventricular non-compaction (LVNC) is a feature of the heart muscle that has been observed in the general population, and reported in conjunction with a DCM phenotype as well as numerous other cardiovascular phenotypes; its relationship (if any) to the presence or severity of the DCM phenotype remains unknown [Hershberger et al 2017, Ross & Semsarian 2018, Ross et al 2020].

Table 2. Nonsyndromic Dilated Cardiomyopathy Genes

Gene ¹	MOI	% of DCM Caused by Pathogenic Variants in Gene ²	ClinGen Gene Validity Classification	Distinguishing Features	Allelic Disorders ³	OMIM Gene Entry
<i>TTN</i> ⁴	AD	15%-20%	Definitive	DCM is assoc w/ truncating variants	LGMD2J; hereditary myopathy w/early respiratory failure; Udd distal myopathy-tibial muscular dystrophy; Salih myopathy; HCM	188840
<i>LMNA</i>	AD	6%	Definitive	Arrhythmia & conduction system disease	Selected examples: partial lipodystrophy; CMT2B1 (See CMT Overview .); Emery-Dreifuss muscular dystrophy; Hutchinson-Gilford progeria syndrome	150330

Table 2. continued from previous page.

Gene ¹	MOI	% of DCM Caused by Pathogenic Variants in Gene ²	ClinGen Gene Validity Classification	Distinguishing Features	Allelic Disorders ³	OMIM Gene Entry
<i>MYH7</i>	AD	4%	Definitive		Laing distal myopathy; HCM; myosin storage myopathy; scapuloperoneal myopathy	160760
<i>FLNC</i>	AD	2%-4%	Definitive	Arrhythmia & conduction system disease	Myofibrillar myopathy; HCM; RCM; distal myopathy	102565
<i>BAG3</i>	AD	3%	Definitive		Myofibrillar myopathy	603883
<i>TNNT2</i>	AD	3%	Definitive		HCM; RCM	191045
<i>RBM20</i> ⁵	AD	2%	Definitive	Arrhythmia & conduction system disease; DCM is assoc w/hot spot in exon 9.		613171
<i>SCN5A</i>	AD	2%	Definitive	Arrhythmia & conduction system disease	Selected examples: Long QT syndrome; Brugada syndrome; idiopathic ventricular fibrillation	600163
<i>DES</i>	AD	<1%	Definitive	Arrhythmia & neuromuscular involvement	Myofibrillar myopathy; neurogenic scapuloperoneal syndrome, Kaeser type	125660
<i>PLN</i>	AD	<1%	Definitive	Arrhythmia & conduction system disease	HCM	172405
<i>TNNC1</i>	AD	<1%	Definitive		HCM	191040
<i>DSP</i>	AD	Unknown	Strong	Arrhythmia & conduction system disease; possible right ventricular involvement	Arrhythmogenic right ventricular cardiomyopathy; epidermolysis bullosa, lethal acantholytic; keratosis palmoplantaris striata II; skin fragility-woolly hair syndrome; Carvajal syndrome	125647
<i>ACTC1</i>	AD	<1%	Moderate		HCM; atrial septal defect	102540
<i>ACTN2</i>	AD	<1%	Moderate		HCM; myopathy	102573
<i>TPM1</i>	AD	<1%	Moderate		HCM	191010
<i>JPH2</i> ⁶	AD AR	Unknown	Moderate		HCM	605267
<i>NEXN</i>	AD	Unknown	Moderate		HCM	613121
<i>TNNI3</i>	AD AR	<1%	Moderate		HCM; RCM	191044
<i>VCL</i>	AD	Unknown	Moderate		HCM	193065
<i>MYH6</i>	AD	4%	Limited		HCM; atrial septal defect	160710

Table 2. continued from previous page.

Gene ¹	MOI	% of DCM Caused by Pathogenic Variants in Gene ²	ClinGen Gene Validity Classification	Distinguishing Features	Allelic Disorders ³	OMIM Gene Entry
<i>MYPN</i>	AD	3%	Limited		RCM; HCM; nemaline myopathy	608517
<i>MYBPC3</i>	AD	2%	Limited		HCM	600958
<i>CSRP3</i>	AD	<1%	Limited		HCM	600824
<i>ILK</i>	AD	<1%	Limited			602366
<i>LAMA4</i>	AD	<1%	Limited			600133
<i>LDB3</i>	AD	<1%	Limited		HCM; myofibrillar myopathy	605906, PS601419
<i>PSEN2</i>	AD	<1%	Limited		Early-onset Alzheimer disease	600759
<i>SGCD</i>	AD	<1%	Limited		LGMD2F / LGMDR6	601411
<i>TCAP</i>	AD	<1%	Limited		HCM; LGMD2G / LGMDR7	604488
<i>ABCC9</i>	AD	Unknown	Limited		Familial atrial fibrillation; Cantú syndrome (hypertrichotic osteochondrodysplasia)	601439
<i>ANKRD1</i>	AD	Unknown	Limited			609599
<i>CTF1</i>	AD	Unknown	Limited			600435
<i>DSG2</i>	AD	Unknown	Limited	Possible right ventricular involvement	Arrhythmogenic right ventricular cardiomyopathy	125671
<i>DTNA</i>	AD	Unknown	Limited		Congenital heart defects	601239
<i>EYA4</i>	AD	Unknown	Limited	Hearing loss	<i>DFNA10</i> nonsyndromic hearing loss & deafness (See Hereditary Hearing Loss and Deafness Overview.)	603550
<i>GATAD1</i>	AR	Unknown	Limited			614518
<i>MYL2</i>	AD	Unknown	Limited		HCM	160781
<i>NEBL</i>	AD	Unknown	Limited			605491
<i>NKX2-5</i>	AD	Unknown	Limited		Congenital heart defects	600584
<i>OBSCN</i>	AD	Unknown	Limited			608616
<i>PLEKHM2</i>	AR	Unknown	Limited			609613
<i>PRDM16</i>	AD	Unknown	Limited			605557
<i>TBX20</i>	AD	Unknown	Limited		Atrial septal defect	606061

Table 2. continued from previous page.

Gene ¹	MOI	% of DCM Caused by Pathogenic Variants in Gene ²	ClinGen Gene Validity Classification	Distinguishing Features	Allelic Disorders ³	OMIM Gene Entry
<i>TNNI3K</i>	AD	Unknown	Limited	Arrhythmia & conduction system disease	Cardiac conduction disease	613932

See [Dilated Cardiomyopathy: OMIM Phenotypic Series](#) to view genes associated with this phenotype in OMIM.

AD = autosomal dominant; AR = autosomal recessive; CMT = Charcot-Marie-Tooth hereditary neuropathy; HCM = [hypertrophic cardiomyopathy](#); LGMD = limb-girdle muscular dystrophy; LGMDR = limb-girdle muscular dystrophy autosomal recessive; MOI = mode of inheritance; RCM = restrictive cardiomyopathy; XL = X-linked

1. Genes are organized first by strength of ClinGen classification, then frequency of causation of DCM, and then alphabetically.

2. The percentages provided (based on ≥ 2 reports screening large numbers of probands with HNDCM) should be interpreted as preliminary estimates.

3. Allelic disorders = other phenotypes caused by pathogenic variants in the same gene.

4. Note: Although 10%-20% of DCM in three cohorts (with or without a family history of DCM) was attributed to *TTN* pathogenic truncating variants [Herman et al 2012], determining the role of pathogenic variants in *TTN* in DCM is difficult given that: (a) 3% of controls also have truncating variants; and (b) *TTN* pathogenic truncating variants have not segregated with DCM in all families with DCM [Norton et al 2013]. Truncating *TTN* variants found in individuals with DCM have been reported to cluster in the A-band region of titin, the protein encoded by *TTN* [Roberts et al 2015]. To date, *TTN* missense variants have not been associated with disease.

5. The hot spot for pathogenic and likely pathogenic variants associated with DCM is located in exon 9 of *RBM20*; it is unclear if non-hot spot variants in *RBM20* can be associated with DCM.

6. Sabater-Molina et al [2016], Miura et al [2020]

3. Establishing (When Possible) the Specific Genetic Cause of DCM

Molecular genetic testing should be offered to every individual of any age with nonischemic DCM [Hershberger et al 2018] including those with peripartum or pregnancy-associated cardiomyopathy (PPCM/PACM) [Goli et al 2021] (Figure 1). See Table 2 for a current list of known DCM-related genes. The purpose of establishing a molecular diagnosis of DCM is to inform risk assessment of relatives of a proband (Section 4).

Variants (pathogenic, likely pathogenic, or of unknown significance) in more than 30 genes have been identified in up to 30%-35% of individuals with familial DCM (i.e., in ≥ 2 first-degree family members) [Hershberger et al 2013, Jordan & Hershberger 2021] or in simplex cases (i.e., in only 1 family member) [Hershberger et al 2008, Hershberger et al 2010b, Pugh et al 2014, Morales et al 2020]. The detection rate of pathogenic and likely pathogenic variants is about 27% [Pugh et al 2014].

A cardiomyopathy multigene panel that includes the genes with a ClinGen classification of definitive, strong, or moderate (as listed in Table 2) is most likely to identify the genetic cause of the condition while also limiting identification of variants of uncertain significance and pathogenic variants in genes that do not explain the underlying phenotype.

Note: (1) The genes included in the panel and the diagnostic sensitivity of the testing used for each gene vary by laboratory and are likely to change over time. (2) Some multigene panels may include genes not associated with the condition discussed in this *GeneReview*. (3) In some laboratories, panel options may include a custom laboratory-designed panel and/or custom phenotype-focused exome analysis that includes genes specified by the clinician. (4) Methods used in a panel may include sequence analysis, deletion/duplication analysis, and/or other non-sequencing-based tests.

For an introduction to multigene panels click [here](#). More detailed information for clinicians ordering genetic tests can be found [here](#).

Health care providers ordering genetic testing should be familiar with the genetics of DCM [Burkett & Hershberger 2005, Judge 2009, Caleshu et al 2010, Hershberger et al 2010a, Hershberger & Siegfried 2011, Hershberger et al 2013, Jordan et al 2021]. Given the complexity of interpreting genetic test results and their implications for surveillance and management, health care providers should consider referral to a cardiovascular genetics center or a genetic counselor specializing in cardiac genetics (see [NSGC - Find a Genetic Counselor](#) or [ABGC Find a Certified Genetic Counselor](#) search tools).

4. Genetic Risk Assessment and Cardiac Surveillance of At-Risk Relatives for Detection of Early Treatable Manifestations of DCM

Cardiovascular screening of asymptomatic first-degree family members of an individual with DCM can allow early detection of DCM, prompt initiation of treatment, and improvement in long-term outcome [Morales & Hershberger 2015]. Clarification of the genetic status of first-degree family members of an individual with DCM can inform who is at risk and the recommended frequency of subsequent cardiovascular screening [Hershberger et al 2018].

A basic view of nonsyndromic dilated cardiomyopathy (DCM) genetic risk assessment and cardiac surveillance for at-risk relatives is presented in this section; issues that may be specific to a given family or genetic cause of nonsyndromic DCM are not comprehensively addressed.

Note: If a proband has a specific syndrome associated with DCM (e.g., [Barth syndrome](#) or [Duchenne muscular dystrophy](#)), counseling for that condition is indicated (see Table 1). Genetic risk assessment in families with syndromic DCM is not discussed further in this section.

Genetic Risk Assessment

Genetic counseling is the process of providing individuals and families with information on the nature, mode(s) of inheritance, and implications of genetic disorders to help them make informed medical and personal decisions. The following section deals with genetic risk assessment and the use of family history and genetic testing to clarify genetic status for family members; it is not meant to address all personal, cultural, or ethical issues that may arise or to substitute for consultation with a genetics professional. —ED.

Nonsyndromic dilated cardiomyopathy (DCM) is typically inherited in an autosomal dominant manner. *JPH2*- and *TNNI3*-related nonsyndromic DCM can be inherited in an autosomal dominant or autosomal recessive manner.

DMD-related DCM – which can present as syndromic DCM in males or isolated DCM in carrier females (Table 1) – is inherited in an X-linked manner (see [Dystrophinopathies](#)).

Autosomal Dominant Inheritance – Risk to Family Members

Parents of a proband

- Some individuals diagnosed as having autosomal dominant DCM have an affected parent.
- Some individuals diagnosed with autosomal dominant DCM have the disorder as the result of a *de novo* pathogenic variant. The proportion of individuals with autosomal dominant DCM caused by a *de novo* pathogenic variant is unknown.
- When evaluating for autosomal dominant inheritance, both the maternal and paternal lineages should be considered as possibly contributing to familial DCM. In an unknown proportion of cases, both parents

may have evidence of DCM and/or DCM-related pathogenic variants, and thus the proband may have inherited pathogenic variants from one or both parents [Liu et al 2015, Cowan et al 2018].

- Molecular genetic testing is recommended for the parents of the proband to confirm their genetic status.
- If the pathogenic variant identified in the proband is not identified in either parent and parental identity testing has confirmed biological maternity and paternity, the following possibilities should be considered:
 - The proband has a *de novo* pathogenic variant.
 - The proband inherited a pathogenic variant from a parent with germline (or somatic and germline) mosaicism. Note: Testing of parental leukocyte DNA may not detect all instances of somatic mosaicism and will not detect a pathogenic variant that is present in the germ cells only.

Sibs of a proband. The risk to the sibs of the proband depends on the clinical/genetic status of the proband's parents:

- If a parent of the proband is affected and/or has a DCM-related pathogenic or likely pathogenic variant, the risk to the sibs of inheriting the variant is 50%. Because of variable expression and reduced penetrance, no predictions can be made regarding age of onset or severity of disease.
- If both parents of a proband have a DCM-related pathogenic or likely pathogenic variant, sibs have a 75% chance of inheriting one or two DCM-related variants and a 25% chance of inheriting neither pathogenic variant.
- If the parents are clinically unaffected but their genetic status is unknown, sibs are still presumed to be at increased risk for DCM because of the possibility of reduced penetrance in a heterozygous parent or parental germline mosaicism.

Offspring of a proband. Each child of an individual with autosomal dominant DCM has a 50% chance of inheriting the DCM-related pathogenic variant.

Autosomal Recessive Inheritance – Risk to Family Members

Parents of a proband

- The parents of an affected individual are obligate heterozygotes (i.e., presumed to have one *JPH2* or *TNNI3* pathogenic variant based on family history).
- Molecular genetic testing is recommended for the parents of a proband to confirm that both parents are heterozygous for a *JPH2* or *TNNI3* pathogenic variant and to allow reliable recurrence risk assessment. If a pathogenic variant is detected in only one parent and parental identity testing has confirmed biological maternity and paternity, the following possibilities should be considered:
 - One of the pathogenic variants identified in the proband occurred as a *de novo* event in the proband or as a postzygotic *de novo* event in a mosaic parent [Jónsson et al 2017].
 - Uniparental isodisomy for the parental chromosome with the pathogenic variant resulted in homozygosity for the pathogenic variant in the proband.
- The heterozygous parents of a child with biallelic pathogenic variants in *JPH2* or *TNNI3* may be at risk for DCM. *JPH2*-related DCM and *TNNI3*-related DCM can be inherited in an autosomal recessive or an autosomal dominant manner and the mechanisms to distinguish between pathogenic variants that solely incur risk in a dominant or recessive model are currently unknown. Therefore, it is reasonable for parents of the proband to undergo cardiac surveillance even if they have only one of the two pathogenic variants identified in the affected proband.

Sibs of a proband

- If both parents are known to be heterozygous for a *JPH2* or *TNNI3* pathogenic variant, each sib of an affected individual has at conception a 25% chance of inheriting biallelic pathogenic variants, a 50%

chance of inheriting one of the two pathogenic variants identified in the proband, and a 25% chance of inheriting neither pathogenic variant.

- A sib who is heterozygous for one of the pathogenic variants identified in the proband may be at risk for DCM. *JPH2* and *TNNI3* pathogenic variants that solely incur risk in a dominant or recessive model are not distinguishable at this time. Therefore, it is reasonable for sibs of the proband to undergo cardiac surveillance even if they have only one of the two pathogenic variants identified in the affected proband.

Offspring of a proband. The offspring of an individual with autosomal recessive *JPH2*- or *TNNI3*-related DCM are obligate heterozygotes for a pathogenic variant in *JPH2* or *TNNI3* and may be at risk for DCM. Offspring should therefore undergo cardiac surveillance for DCM.

Cardiac Surveillance

It is appropriate to clarify the clinical and genetic status of asymptomatic family members at risk for DCM prior to the onset of manifestations to identify those with asymptomatic DCM and permit initiation of medical therapy aimed at preventing/delaying the morbidity of late-stage symptomatic disease [Morales & Hershberger 2015].

The following recommendations for surveillance of asymptomatic at-risk family members reflect the practice guidelines of the Heart Failure Society of America [Hershberger et al 2018].

If the Proband Has a Known Pathogenic Variant (or Pathogenic Variants) in a DCM-Related Gene

See Figure 1, teal box ("Pathogenic variant identified"). Molecular genetic testing is recommended for parents, sibs, offspring, and other at-risk family members in order to clarify their genetic status.

Those identified as having a familial DCM-related pathogenic variant have an increased lifetime risk for DCM and, when asymptomatic, should undergo cardiovascular clinical screening at intervals based on the individual's age [Hershberger et al 2018].

Note: Asymptomatic at-risk relatives who do not meet criteria for DCM (with other causes ruled out) may represent early DCM when echocardiogram results are ambiguous (e.g., left ventricular enlargement with normal systolic function, decreased ejection fraction but normal-sized left ventricle) and/or echocardiogram results are normal but EKG results are abnormal (e.g., significant conduction system disease and/or arrhythmias).

In general, family members without the DCM-related pathogenic variant identified in the proband are no longer considered to be at increased risk for DCM and thus may be discharged from cardiac surveillance. However, because multiple variants in DCM-associated genes have been observed in individuals with nonsyndromic DCM [Morales et al 2020] and because families may segregate pathogenic variants in more than one DCM-related gene [Liu et al 2015, Cowan et al 2018], thorough individualized risk assessment through clinical, genetic, and family history analysis is warranted to determine if discharge from high-risk cardiac surveillance is appropriate.

If the Specific Genetic Cause of DCM in the Proband Has Not Been Identified

See Figure 1, orange box ("Pathogenic variant not identified"). Perform cardiovascular screening on asymptomatic at-risk family members at intervals based on the individual's age [Hershberger et al 2018].

Note: Asymptomatic at-risk relatives who do not meet criteria for DCM (with other causes ruled out) may represent early DCM when echocardiogram results are ambiguous (e.g., left ventricular enlargement with normal systolic function, decreased ejection fraction but normal-sized left ventricle) and/or echocardiogram

results are normal but EKG results are abnormal (e.g., significant conduction system disease and/or arrhythmias).

If a first-degree at-risk relative shows evidence of dilated cardiomyopathy, a diagnosis of familial DCM is made and the surveillance recommendations should extend to that person's first-degree relatives.

Future additional genetic testing for the proband (and other informative family members) may be considered when:

- Multigene panels are expanded to include more genes and test sensitivity increases (e.g., resulting from better coverage of the genes included and improved detection of deletions/duplications); and
- Genomic testing (exome sequencing and genome sequencing) becomes more suitable for clinical use.

Resources

GeneReviews staff has selected the following disease-specific and/or umbrella support organizations and/or registries for the benefit of individuals with this disorder and their families. GeneReviews is not responsible for the information provided by other organizations. For information on selection criteria, click [here](#).

- **American Heart Association**
[Dilated Cardiomyopathy \(DCM\)](#)
- **Cardiomyopathy UK**
United Kingdom
Phone: 0800 018 1024 (UK only)
Email: contact@cardiomyopathy.org
www.cardiomyopathy.org
- **Children's Cardiomyopathy Foundation**
Phone: 866-808-2873 (toll-free)
Fax: 201-227-7016
Email: info@childrenscardiomyopathy.org
www.childrenscardiomyopathy.org
- **DCM Foundation**
Phone: 833-DCM-HOPE (833-326-4673)
Email: Info@DCMFoundation.org
www.dcmfoundation.org
- **MedlinePlus**
[Familial dilated cardiomyopathy](#)
- **Dilated Cardiomyopathy Research Project**
Phone: 877-800-8430
Email: DCM.Research@osumc.edu
www.dcmproject.com

Chapter Notes

Author Notes

Web: [Dilated Cardiomyopathy Research Project](#)

The Dilated Cardiomyopathy Research Project, originally launched in 1993 by Dr Ray Hershberger while at the Oregon Health & Science University, aims to advance our understanding of dilated cardiomyopathy genetics. Multiple studies including data from more than 1500 families affected by DCM across the country have contributed genetic and clinical information to this research program. This multi-institutional effort, still led by Dr Hershberger and now housed at The Ohio State University, leverages the many sites collaborating in the DCM Consortium to identify families eligible for studies within the DCM Research Project, including the recently completed DCM Precision Medicine Study and the ongoing DCM Discovery Study. More information about the DCM Project, affiliated research studies, and other information and resources can be found on the website (www.dcmproject.com) or by emailing dcm.research@osumc.edu.

Acknowledgments

We are deeply grateful for the numerous families who have contributed to the research efforts in the DCM Research Project for nearly three decades. Without their gracious participation and engagement, the advances driven by the work of the DCM Research Project to the field would not be possible.

Author History

Ray E Hershberger, MD (2007-present)

Elizabeth Jordan, MS, LGC (2021-present)

Jessica D Kushner, MS, CGC; Oregon Health & Science University (2007-2013)

Ana Morales, MS, CGC; The Ohio State University (2013-2021)

Sharie Parks, PhD; Oregon Health & Science University (2007-2013)

Revision History

- 7 April 2022 (aa) Revision: Cardiac surveillance recommendations (If the Proband Has a Known Pathogenic Variant[s] in a DCM-Related Gene)
- 29 July 2021 (ha) Comprehensive update posted live
- 23 August 2018 (bp) Comprehensive update posted live
- 24 September 2015 (me) Comprehensive update posted live
- 9 May 2013 (me) Comprehensive update posted live
- 19 March 2009 (cd) Revision: sequence analysis and prenatal testing available clinically for TCAP, ABCC9, VCL, ACTN2, and CSRP3
- 10 July 2008 (cd) Revision: clinical testing available for TTN mutations as a cause of dilated cardiomyopathy
- 27 July 2007 (me) Review posted live
- 6 December 2006 (jdk) Original submission

References

Published Guidelines / Consensus Statements

- Hershberger RE, Givertz MM, Ho CY, Judge DP, Kantor PF, McBride KL, Morales A, Taylor MRG, Vatta M, Ware SM, et al. Genetic evaluation of cardiomyopathy: a clinical practice resource of the American College of Medical Genetics and Genomics (ACMG). *Genet Med.* 2018;20:899-909. [[PubMed](#)]
- Committee on Bioethics, Committee on Genetics, and American College of Medical Genetics and Genomics Social, Ethical, Legal Issues Committee. Ethical and policy issues in genetic testing and screening of children. Available [online](#). 2013. Accessed 8-11-22.
- National Society of Genetic Counselors. Position statement on genetic testing of minors for adult-onset conditions. Available [online](#). 2018. Accessed 8-11-22.

Literature Cited

- Burkett EL, Hershberger RE. Clinical and genetic issues in familial dilated cardiomyopathy. *J Am Coll Cardiol*. 2005;45:969–81. PubMed PMID: 15808750.
- Caleshu C, Day S, Rehm HL, Baxter S. Use and interpretation of genetic tests in cardiovascular genetics. *Heart*. 2010;96:1669–75. PubMed PMID: 20937756.
- Cowan JR, Kinnamon DD, Morales A, Salyer L, Nickerson DA, Hershberger RE. Multigenic disease and bilineal inheritance in dilated cardiomyopathy is illustrated in nonsegregating LMNA pedigrees. *Circ Genom Precis Med*. 2018;11:e002038. PubMed PMID: 30012837.
- Dellefave L, McNally EM. The genetics of dilated cardiomyopathy. *Curr Opin Cardiol*. 2010;25:198–204. PubMed PMID: 20186049.
- Garcia-Pavia P, Kim Y, Restrepo-Cordoba MA, Lunde IG, Wakimoto H, Smith AM, Toepfer CN, Getz K, Gorham J, Patel P, Ito K, Willcox JA, Arany Z, Li J, Owens AT, Govind R, Nuñez B, Mazaika E, Bayes-Genis A, Walsh R, Finkelman B, Lupon J, Whiffin N, Serrano I, Midwinter W, Wilk A, Bardaji A, Ingold N, Buchan R, Tayal U, Pascual-Figal DA, de Marvao A, Ahmad M, Garcia-Pinilla JM, Pantazis A, Dominguez F, John Baksi A, O'Regan DP, Rosen SD, Prasad SK, Lara-Pezzi E, Provencio M, Lyon AR, Alonso-Pulpon L, Cook SA, DePalma SR, Barton PJR, Aplenc R, Seidman JG, Ky B, Ware JS, Seidman CE. Genetic variants associated with cancer therapy-induced cardiomyopathy. *Circulation*. 2019;140:31–41. PubMed PMID: 30987448.
- Goli R, Li J, Brandimarto J, Levine LD, Riis V, McAfee Q, DePalma S, Haghghi A, Seidman JG, Seidman CE, Jacoby D, Macones G, Judge DP, Rana S, Margulies KB, Cappola TP, Alharethi R, Damp J, Hsich E, Elkayam U, Sheppard R, Alexis JD, Boehmer J, Kamiya C, Gustafsson F, Damm P, Ersbøll AS, Goland S, Hilfiker-Kleiner D, McNamara DM, Arany Z, et al. Genetic and phenotypic landscape of peripartum cardiomyopathy. *Circulation*. 2021;143:1852–62. PubMed PMID: 33874732.
- Herman DS, Lam L, Taylor MR, Wang L, Teekakirikul P, Christodoulou D, Conner L, DePalma SR, McDonough B, Sparks E, Teodorescu DL, Cirino AL, Banner NR, Pennell DJ, Graw S, Merlo M, Di Lenarda A, Sinagra G, Bos JM, Ackerman MJ, Mitchell RN, Murry CE, Lakdawala NK, Ho CY, Barton PJ, Cook SA, Mestroni L, Seidman JG, Seidman CE. Truncations of titin causing dilated cardiomyopathy. *N Engl J Med*. 2012;366:619–28. PubMed PMID: 22335739.
- Hershberger RE, Cowan J, Morales A, Siegfried JD. Progress with genetic cardiomyopathies: screening, counseling, and testing in dilated, hypertrophic, and arrhythmogenic right ventricular dysplasia/cardiomyopathy. *Circ Heart Fail*. 2009;2:253–61. PubMed PMID: 19808347.
- Hershberger RE, Givertz MM, Ho CY, Judge DP, Kantor PF, McBride KL, Morales A, Taylor MRG, Vatta M, Ware SM. Genetic evaluation of cardiomyopathy—a Heart Failure Society of America Practice Guideline. *J Card Fail*. 2018;24:281–302. PubMed PMID: 29567486.
- Hershberger RE, Hedges DJ, Morales A. Dilated cardiomyopathy: the complexity of a diverse genetic architecture. *Nat Rev Cardiol*. 2013;10:531–47. PubMed PMID: 23900355.
- Hershberger RE, Morales A, Cowan J. Is left ventricular noncompaction a trait, phenotype, or disease? The evidence points to phenotype. *Circ Cardiovasc Genet*. 2017;10:e001968. PubMed PMID: 29212902.
- Hershberger RE, Morales A, Siegfried JD. Clinical and genetic issues in dilated cardiomyopathy: A review for genetics professionals. *Genet Med*. 2010a;12:655–67. PubMed PMID: 20864896.
- Hershberger RE, Norton N, Morales A, Li D, Siegfried JD, Gonzalez-Quintana J. Coding sequence rare variants identified in MYBPC3, MYH6, TPM1, TNNC1, and TNNI3 from 312 patients with familial or idiopathic dilated cardiomyopathy. *Circ Cardiovasc Genet*. 2010b;3:155–61. PubMed PMID: 20215591.

- Hershberger RE, Parks SB, Kushner JD, Li D, Ludwigsen S, Jakobs P, Nauman D, Burgess D, Partain J, Litt M. Coding sequence mutations identified in MYH7, TNNT2, SCN5A, CSRP3, LBD3, and TCAP from 313 patients with familial or idiopathic dilated cardiomyopathy. *Clin Transl Sci*. 2008;1:21–6. PubMed PMID: 19412328.
- Hershberger RE, Siegfried JD. State of the art review. Update 2011: clinical and genetic issues in familial dilated cardiomyopathy. *J Am Coll Cardiol*. 2011;57:1641–9. PubMed PMID: 21492761.
- Jónsson H, Sulem P, Kehr B, Kristmundsdottir S, Zink F, Hjartarson E, Hardarson MT, Hjorleifsson KE, Eggertsson HP, Gudjonsson SA, Ward LD, Arnadottir GA, Helgason EA, Helgason H, Gylfason A, Jonasdottir A, Jonasdottir A, Rafnar T, Frigge M, Stacey SN, Th Magnusson O, Thorsteinsdottir U, Masson G, Kong A, Halldorsson BV, Helgason A, Gudbjartsson DF, Stefansson K. Parental influence on human germline de novo mutations in 1,548 trios from Iceland. *Nature*. 2017;549:519–22. PubMed PMID: 28959963.
- Jordan E, Hershberger RE. Considering complexity in the genetic evaluation of dilated cardiomyopathy. *Heart*. 2021;107:106–12. PubMed PMID: 33109712.
- Jordan E, Peterson L, Ai T, Asatryan B, Bronicki L, Brown E, Celeghin R, Edwards M, Fan J, Ingles J, James CA, Jarinova O, Johnson R, Judge DP, Lahrouchi N, Lekanne Deprez RH, Lumbers RT, Mazzarotto F, Medeiros Domingo A, Miller RL, Morales A, Murray B, Peters S, Pilichou K, Protonotarios A, Semsarian C, Shah P, Syrris P, Thaxton C, van Tintelen JP, Walsh R, Wang J, Ware J, Hershberger RE. An evidence-based assessment of genes in dilated cardiomyopathy. *Circulation*. 2021;144:7–19. PubMed PMID: 33947203.
- Judge DP. Use of genetics in the clinical evaluation of cardiomyopathy. *JAMA*. 2009;302:2471–6. PubMed PMID: 19996403.
- Liu GS, Morales A, Vafiadaki E, Lam CK, Cai WF, Haghghi K, Adly G, Hershberger RE, Kranias EG. A novel human R25C-phospholamban mutation is associated with super-inhibition of calcium cycling and ventricular arrhythmia. *Cardiovasc Res*. 2015;107:164–74. PubMed PMID: 25852082.
- Miura A, Kondo H, Yamamoto T, Okumura Y, Nishio H. Sudden unexpected death of infantile dilated cardiomyopathy with JPH2 and PKD1 gene variants. *Int Heart J*. 2020;61:1079–83. PubMed PMID: 32879264.
- Morales A, Hershberger RE. The rationale and timing of molecular genetic testing for dilated cardiomyopathy. *Can J Cardiol*. 2015;31:1309–12. PubMed PMID: 26518443.
- Morales A, Kinnamon DK, Jordan E, Platt J, Vatta M, Dorschner MO, Starkey CA, Mead JO, Ai T, Burke W, Gastier-Foster J, Jarvik GP, Rehm HL, Nickerson DA, Hershberger RE, et al. Variant interpretation for dilated cardiomyopathy: refinement of the American College of Medical Genetics and Genomics/ClinGen guidelines for the DCM Precision Medicine Study. *Circ Genom Precis Med*. 2020;13:e002480. PubMed PMID: 32160020.
- Norton N, Li D, Rampersaud E, Morales A, Martin ER, Zuchner S, Guo S, Gonzalez M, Hedges DJ, Robertson PD, Krumm N, Nickerson DA, Hershberger RE, et al. Exome sequencing and genome-wide linkage analysis in 17 families illustrate the complex contribution of TTN truncating variants to dilated cardiomyopathy. *Circ Cardiovasc Genet*. 2013;6:144–53. PubMed PMID: 23418287.
- Pugh TJ, Kelly MA, Gowrisankar S, Hynes E, Seidman MA, Baxter SM, Bowser M, Harrison B, Aaron D, Mahanta LM, Lakdawala NK, McDermott G, White ET, Rehm HL, Lebo M, Funke BH. The landscape of genetic variation in dilated cardiomyopathy as surveyed by clinical DNA sequencing. *Genet Med*. 2014;16:601–8. PubMed PMID: 24503780.
- Roberts AM, Ware JS, Herman DS, Schafer S, Baksi J, Bick AG, Buchan RJ, Walsh R, John S, Wilkinson S, Mazzarotto F, Felkin LE, Gong S, MacArthur JA, Cunningham F, Flannick J, Gabriel SB, Altshuler DM, Macdonald PS, Heinig M, Keogh AM, Hayward CS, Banner NR, Pennell DJ, O'Regan DP, San TR, de Marvao A, Dawes TJ, Gulati A, Birks EJ, Yacoub MH, Radke M, Gotthardt M, Wilson JG, O'Donnell CJ, Prasad SK,

Barton PJ, Fatkin D, Hubner N, Seidman JG, Seidman CE, Cook SA. Integrated allelic, transcriptional, and phenomic dissection of the cardiac effects of titin truncations in health and disease. *Sci Transl Med*. 2015;7:270ra6.

Ross SB, Jones K, Blanch B, Puranik R, McGeechan K, Barratt A, Semsarian C. A systematic review and meta-analysis of the prevalence of left ventricular non-compaction in adults. *Eur Heart J*. 2020;41:1428–36. PubMed PMID: 31143950.

Ross SB, Semsarian C. Clinical and genetic complexities of left ventricular noncompaction: preventing overdiagnosis in a disease we do not understand. *JAMA Cardiol*. 2018;3:1033–4. PubMed PMID: 30140926.

Sabater-Molina M, Navarro M, García-Molina Sáez E, Garrido I, Pascual-Figal D, González Carrillo J, Gimeno Blanes JR. Mutation in JPH2 cause dilated cardiomyopathy. *Clin Genet*. 2016;90:468–9. PubMed PMID: 27471098.

Sen-Chowdhry S, Syrris P, Prasad SK, Hughes SE, Merrifield R, Ward D, Pennell DJ, McKenna WJ. Left-dominant arrhythmogenic cardiomyopathy: an under-recognized clinical entity. *J Am Coll Cardiol*. 2008;52:2175–87. PubMed PMID: 19095136.

Sivasankaran S, Sharland GK, Simpson JM. Dilated cardiomyopathy presenting during fetal life. *Cardiol Young*. 2005;15:409–16. PubMed PMID: 16014190.

License

GeneReviews® chapters are owned by the University of Washington. Permission is hereby granted to reproduce, distribute, and translate copies of content materials for noncommercial research purposes only, provided that (i) credit for source (<http://www.genereviews.org/>) and copyright (© 1993-2024 University of Washington) are included with each copy; (ii) a link to the original material is provided whenever the material is published elsewhere on the Web; and (iii) reproducers, distributors, and/or translators comply with the [GeneReviews® Copyright Notice and Usage Disclaimer](#). No further modifications are allowed. For clarity, excerpts of GeneReviews chapters for use in lab reports and clinic notes are a permitted use.

For more information, see the [GeneReviews® Copyright Notice and Usage Disclaimer](#).

For questions regarding permissions or whether a specified use is allowed, contact: admasst@uw.edu.