

Title: Prolidase Deficiency *GeneReview* - Laboratory techniques used in the diagnosis of prolidase deficiency

Authors: Rossignol F, Wang H, Ferreira C

Date: June 2015; chapter updated 2022

Note: The following information is provided by the authors listed above and has not been reviewed by *GeneReviews* staff.

**Measurement of prolidase enzyme activity in erythrocytes, leukocytes, or cultured fibroblasts.** Assays are performed with high concentrations (10-50 mM) of imidodipeptides as substrates [Hechtman 2014]. The preferred substrate is Gly-Pro [Royce & Steinmann 2002]. Following incubation at 37°C, the reaction is terminated by heating or by adding trichloroacetic acid [Hechtman 2014]. The released proline is quantitated by spectrophotometry, using either Chinard's original method [Chinard 1952] or its modification by Myara et al [1982]. Other techniques used for measuring prolidase activity include reverse-phase high-performance liquid chromatography [Harada et al 1990], capillary electrophoresis [Zanaboni et al 1997], and more recently MALDI-TOF mass spectrometry [Kurien et al 2004].

A review of the different laboratory techniques used in the diagnosis of prolidase deficiency can be found elsewhere [Kurien et al 2006, Viglio et al 2006].

## References

Chinard FP. Photometric estimation of proline and ornithine. *J Biol Chem.* 1952;199:91–5.

Harada M, Fukasawa KM, Hiraoka BY, Fukasawa K, Mogi M. High-performance liquid chromatographic procedure for the determination of serum prolidase activity. *J Chromatogr.* 1990;530:116–21.

Hechtman P. Prolidase deficiency. In: Beaudet AL, Vogelstein B, Kinzler KW, Antonarakis SE, Ballabio A, Gibson KM, Mitchell G, eds. *The Online Metabolic and Molecular Bases of Inherited Disease*. McGraw-Hill; 2014.

Kurien BT, Patel NC, Porter AC, D'Souza A, Miller D, Matsumoto H, Wang H, Scofield RH. Prolidase deficiency and the biochemical assays used in its diagnosis. *Anal Biochem.* 2006;349:165–75.

Kurien BT, Patel NC, Porter AC, Kurono S, Matsumoto H, Wang H, Scofield RH. Determination of prolidase activity using matrix-assisted laser desorption/ionization time-of-flight mass spectrometry. *Anal Biochem.* 2004;331:224–9.

Myara I, Charpentier C, Lemonnier A. Optimal conditions for prolidase assay by proline colorimetric determination: application to iminodipeptiduria. *Clin Chim Acta.* 1982;125:193–205.

Royce P, Steinmann B. Prolidase deficiency. In: Royce P, Steinmann B. *Connective Tissue and Its Heritable Disorders*. Wiley-Liss; 2002:727–38.

Viglio S, Annovazzi L, Conti B, Genta I, Perugini P, Zanone C, Casado B, Cetta G, Iadarola P. The role of emerging techniques in the investigation of prolidase deficiency: from diagnosis to the development of a possible therapeutical approach. *J Chromatogr B Analyt Technol Biomed Life Sci.* 2006;832:1–8.

Zanaboni G, Viglio S, Dyne KM, Grimm R, Valli M, Cetta G, Iadarola P. Direct monitoring of prolidase activity in cultured skin fibroblasts using capillary electrophoresis. *J Chromatogr B Biomed Sci Appl.* 1997;695:77–84.